



REMEMBER

If your child seems hot and irritable following vaccination you can give Calpol either to aid the discomfort or treat a mild fever – see overleaf.

HEALTH VISITOR

For advice regarding non-urgent matters please ring and leave a message on 01494 426709

It can be helpful to tell your child what is going to happen.

All immunisations should be recorded in the Red Book, so do please bring it with you.

Your immunisation nurse is:

Cate Norridge

ADVICE FOLLOWING IMMUNISATIONS



**Dr Jill Deacon
Dr Martin Davis
Dr Lena Poulton
Dr Sarah Rapsey**



HIGHFIELD SURGERY
Highfield Surgery
Highfield Way
Hazlemere, High Wycombe
HP15 7UW

Tel: 01494 813396/813496

Dear Parents,

Congratulate yourself on helping your child receive one of the most important protective measures available, according to the World Health Organisation. These immunisations should be trouble free, but this leaflet will help if you have any specific concerns.

FIRST IMMUNISATIONS

After the 1st and 3rd immunisations which include Meningitis B it is recommended to give 3 doses of infant Calpol 2.5mls, the first dose immediately after the vaccinations and 2 further doses 4-6 hrs apart, this should minimise the possible temperature.

Occasionally the injection site may become red, sore and swollen around the place where the injection has been given, or a small lump may appear. This may take a few weeks to go away, it is not a cause for concern.

For more information about childhood vaccinations visit our website:

www.highfieldsurgeryhazlemere.co.uk

- Go to Information Leaflets and Links tab (left hand side) and
- Follow link - Guide to Immunisations

Or visit the NHS Choices website:

www.nhs.uk

- Search Vaccinations

FOLLOWING MMR IMMUNISATION

Your baby may have just had a measles, mumps and rubella vaccination – this contains three different live viruses which may act at different times on your child and can produce the following side effects:

- 6-10 days after the immunisation, some children may become feverish (this happens to about one in ten children). Some develop a measles-like rash and go off their food as the measles part of the vaccine starts to work.
- Very rarely a child may have a fit caused by the fever and this is called a 'febrile convulsion'. Please contact the surgery or the NHS 111 Service for advice on this. However, if a child who has not been immunised gets measles, they are five times more likely to have a fit.
- Rarely, children may get mumps-like symptoms (fever and swollen glands) about three weeks after their immunisation as the mumps part of the vaccine starts to work.
- Very rarely, children may get a rash of small bruise-like spots in the six weeks after the vaccination. This is usually caused by the measles or rubella parts of the vaccine. If you see spots like these, and they are unwell take your child to the doctor to be checked.
- Fewer than one child in a million develops encephalitis (swelling of the brain) after the MMR vaccine, and there is very little evidence that it is caused by the vaccine. However, if a child catches measles, the chance of developing encephalitis is between one in 200 and one in 5000.

None of the viruses are infectious in any way so the child can mix with people as normal.



If you are worried or anxious about your baby, please do not hesitate to contact the surgery on the telephone number overleaf. If it is out of hours, you may wish to contact the NHS 111 Service for advice.



CONSENT

If neither Mum nor Dad can bring their baby for their vaccinations please can they provide written consent that they are happy for another carer to bring the child.

Example

I, Kate Smith give consent for my son/daughter, Baby Smith, to have the vaccinations due. They will be brought by (name of carer).

Signed and dated